

Crohn's/Ulcerative Colitis enrollment form

Phone: 855-425-4085 Fax: 855-425-4096 ardonhealth.com



Date needed	Medication start date	Ship to: <input type="checkbox"/> Patient <input type="checkbox"/> Physician <input type="checkbox"/> Other:
-------------	-----------------------	--

Patient information					
Patient name	Date of birth	Phone	Alternate phone		
Address	City	State	ZIP		
Gender: <input type="checkbox"/> Male <input type="checkbox"/> Female	Email	Primary language	Height	Weight	

Prescriber information					
Prescriber name	State License #	NPI #	DEA #		
Group or hospital	Address	City	State	ZIP	
Phone	Fax	Contact person name and phone			

Insurance information: If available, please fax a copy of the prescription and insurance card(s) with this form (front and back).

Clinical	
Date of diagnosis	Diagnosis ICD-10 code: <input type="checkbox"/> K50.90 Crohn's Disease <input type="checkbox"/> K51.90 Ulcerative Colitis <input type="checkbox"/> Other (ICD-10 Code) _____
Previous medications:	Current medications:
Does the patient have a latex allergy? <input type="checkbox"/> Yes <input type="checkbox"/> No	Is the patient also taking methotrexate? <input type="checkbox"/> Yes <input type="checkbox"/> No
Does patient have active/serious infection? <input type="checkbox"/> Yes <input type="checkbox"/> No	Allergies:
Does patient have heart failure? <input type="checkbox"/> Yes <input type="checkbox"/> No	Is the patient new to therapy? <input type="checkbox"/> Yes <input type="checkbox"/> No
Has patient had positive TB test? <input type="checkbox"/> Yes <input type="checkbox"/> No <i>If yes, date of last chest x-ray</i> _____	

Prescription information				
Medication	Dose/strength	Directions	Quantity	Refill
<input type="checkbox"/> Cimzia®	<input type="checkbox"/> Starter Kit	<input type="checkbox"/> Induction Dose: Inject 400 mg SUBQ on day 1, 15, and 29	<input type="checkbox"/> 1 Starter Kit = 6 PFS	0
	<input type="checkbox"/> 200 mg/mL Prefilled Syringe <input type="checkbox"/> 200 mg Vial	<input type="checkbox"/> Maintenance Dose: Inject 400 mg SUBQ every 28 days	<input type="checkbox"/> 2 PFS/Vials	
<input type="checkbox"/> Dupixent®	<input type="checkbox"/> 300 mg/2 mL Pen <input type="checkbox"/> 300 mg/2 mL Prefilled Syringe	<input type="checkbox"/> Inject 300 mg SUBQ every 7 days	<input type="checkbox"/> 4 Pens/PFS	
	<input type="checkbox"/> 40 mg/0.8 mL Pen Crohn's Disease, Ulcerative Colitis Starter (6 pens) <input type="checkbox"/> 40 mg/0.8 mL Prefilled Syringe Pediatric Crohn's Disease Starter (6 syringes) <input type="checkbox"/> 40 mg/0.8 mL Prefilled Syringe Pediatric Crohn's Disease Starter (3 syringes)	<input type="checkbox"/> Adult Crohn's/UC and pediatric Crohn's ≥ 40 kg: Inject 160 mg SUBQ on day 1, 80 mg on day 15, then 40 mg every 14 days thereafter starting on day 29 <input type="checkbox"/> Pediatric Crohn's 17 to < 40 kg: Inject 80 mg SUBQ on day 1, 40 mg on day 15, then 20 mg every 14 days starting on day 29 <input type="checkbox"/> Other: _____	<input type="checkbox"/> 1 Kit	0
<input type="checkbox"/> Humira®	<input type="checkbox"/> 40 mg/0.8 mL Pen	<input type="checkbox"/> Inject 40 mg SUBQ every 14 days	<input type="checkbox"/> 2 Pens/PFS	
	<input type="checkbox"/> 40 mg/0.8 mL Prefilled Syringe	<input type="checkbox"/> Inject 40 mg SUBQ every 7 days	<input type="checkbox"/> 4 Pens/PFS	
	<input type="checkbox"/> 20 mg/0.4 mL Prefilled Syringe	<input type="checkbox"/> Other: _____		
	<input type="checkbox"/> 10 mg/0.2 mL Prefilled Syringe			

Physician signature required	
Product substitution permitted <input checked="" type="checkbox"/> _____ Date _____	Dispense as written <input checked="" type="checkbox"/> _____ Date _____

Ancillary supplies and kits will be provided as needed for administration.

The information included in this FAX is intended for the sole use of the individual to whom it is addressed and may contain information that is privileged, confidential and exempt from disclosure under applicable law. If you are not the intended addressee, nor authorized to receive for the intended addressee, you are hereby notified that you may not use, copy, disclose or distribute this information. If you have received this FAX in error, please contact the sender and destroy the entire document.

<input type="checkbox"/> Humira® (Citrate-free)	Starter Dose <input type="checkbox"/> 80 mg/0.8 mL Pen Crohn's Disease, Ulcerative Colitis Starter (3 pens) <input type="checkbox"/> 40 mg/0.4 mL Pen Crohn's Disease, Ulcerative Colitis Starter (6 pens) <input type="checkbox"/> 80 mg/0.8 mL Prefilled Syringe Pediatric Crohn's Disease Starter (3 syringes) <input type="checkbox"/> 80 mg/0.8 mL and 40 mg/0.4 mL Prefilled Syringe Pediatric Crohn's Disease Starter (2 syringes) <input type="checkbox"/> 80 mg/0.8 mL Pen Pediatric Ulcerative Colitis Starter (4 pens) <input type="checkbox"/> 40 mg/0.4 mL Pen Pediatric Ulcerative Colitis (4 pens) <input type="checkbox"/> 40 mg/0.4 mL Prefilled Syringe Pediatric Ulcerative Colitis (4 syringes)	<input type="checkbox"/> Adult Crohn's/UC and pediatric Crohn's \geq 40 kg: Inject 160 mg SUBQ on day 1, 80 mg on day 15, then 40 mg every 14 days thereafter starting on day 29 <input type="checkbox"/> Pediatric Crohn's 17 to < 40 kg: Inject 80 mg SUBQ on day 1, 40 mg on day 15, then 20 mg every 14 days starting on day 29 <input type="checkbox"/> Pediatric UC \geq 40 kg: Inject 160 mg SUBQ on day 1, 80 mg on day 8, 80 mg on day 15, then begin maintenance dosing starting on day 29 <input type="checkbox"/> Pediatric UC 20 kg to < 40 kg: Inject 80 mg SUBQ on day 1, 40 mg on day 8, 40 mg on day 15, then begin maintenance dosing starting on day 29 <input type="checkbox"/> Other: _____	<input type="checkbox"/> 1 Kit	0
	Maintenance Dose <input type="checkbox"/> 80 mg/0.8 mL CF Pen <input type="checkbox"/> 40 mg/0.4 mL CF Pen <input type="checkbox"/> 40 mg/0.4 mL CF Prefilled Syringe <input type="checkbox"/> 20 mg/0.2 mL CF Prefilled Syringe <input type="checkbox"/> 10 mg/0.1 mL CF Prefilled Syringe	<input type="checkbox"/> Inject 40 mg SUBQ every 14 days <input type="checkbox"/> Inject 80 mg SUBQ every 14 days <input type="checkbox"/> Inject 40 mg SQ every 7 days <input type="checkbox"/> Other: _____	<input type="checkbox"/> 2 Pens/PFS <input type="checkbox"/> 4 Pens/PFS	
<input type="checkbox"/> Omvoh®	<input type="checkbox"/> 100 mg/mL Pen <input type="checkbox"/> 100 mg/mL Prefilled Syringe	<input type="checkbox"/> Maintenance Dose: Inject 200 mg SUBQ at Week 12, followed by every 4 weeks thereafter Has the patient received the IV induction doses already? <input type="checkbox"/> Yes <input type="checkbox"/> No <i>If yes, date of IV induction doses:</i> Week 0: _____ Week 4: _____ Week 8: _____	<input type="checkbox"/> 2 Pens/PFS	
<input type="checkbox"/> Remicade®	<input type="checkbox"/> 100 mg Vial	<input type="checkbox"/> Induction: Infuse _____ mg IV at weeks 0, 2, and 6	_____ vial(s)	0
		<input type="checkbox"/> Maintenance: Infuse _____ mg IV every 8 weeks	_____ vial(s)	
<input type="checkbox"/> Rinvoq®	<input type="checkbox"/> 45 mg XR Tablet	<input type="checkbox"/> Take 1 tablet by mouth once daily	<input type="checkbox"/> 28 Tablets	
	<input type="checkbox"/> 15 mg XR Tablet <input type="checkbox"/> 30 mg XR Tablet	<input type="checkbox"/> Take 1 tablet by mouth once daily	<input type="checkbox"/> 30 Tablets	
<input type="checkbox"/> Simponi®	<input type="checkbox"/> 100 mg/mL Pen <input type="checkbox"/> 100 mg/mL Prefilled Syringe	<input type="checkbox"/> Induction Dose: Inject 200 mg SUBQ day 1, then 100 mg on day 15, then 100 mg every 28 days thereafter	<input type="checkbox"/> 3 Pens/PFS	0
		<input type="checkbox"/> Maintenance Dose: Inject 100 mg SUBQ every 28 days	<input type="checkbox"/> 1 Pen/PFS	
<input type="checkbox"/> Skyrizi®	<input type="checkbox"/> 180 mg/1.2 mL prefilled cartridge with on-body injector	<input type="checkbox"/> Maintenance Dose: Inject 180 mg SUBQ at week 12, followed by every 8 weeks thereafter	<input type="checkbox"/> 1 Kit	
	<input type="checkbox"/> 360 mg/2.4 mL prefilled cartridge with on-body injector	<input type="checkbox"/> Maintenance Dose: Inject 360 mg SUBQ at week 12, followed by every 8 weeks thereafter Has the patient received the IV induction doses already? <input type="checkbox"/> Yes <input type="checkbox"/> No <i>If yes, date of IV induction doses:</i> Week 0: _____ Week 4: _____ Week 8: _____		
<input type="checkbox"/> Stelara®	<input type="checkbox"/> 90 mg Prefilled Syringe	<input type="checkbox"/> Maintenance Dose: Inject 90 mg SUBQ 8 weeks after initial IV dose, followed by every 8 weeks thereafter Has the patient received the IV induction dose already? <input type="checkbox"/> Yes <input type="checkbox"/> No <i>If yes, date of IV induction dose:</i> _____	<input type="checkbox"/> 1 PFS	
<input type="checkbox"/> Velsipity®	<input type="checkbox"/> 2 mg Tablet	<input type="checkbox"/> Take 1 tablet by mouth once daily	<input type="checkbox"/> 30	

Physician signature required

Product substitution permitted

X _____ Date _____

Dispense as written

X _____ Date _____

Ancillary supplies and kits will be provided as needed for administration.

The information included in this FAX is intended for the sole use of the individual to whom it is addressed and may contain information that is privileged, confidential and exempt from disclosure under applicable law. If you are not the intended addressee, nor authorized to receive for the intended addressee, you are hereby notified that you may not use, copy, disclose or distribute this information. If you have received this FAX in error, please contact the sender and destroy the entire document.

<input type="checkbox"/> Xeljanz®	<input type="checkbox"/> 10 mg Tablet	<input type="checkbox"/> Take 1 tablet by mouth 2 times daily	<input type="checkbox"/> 60 Tablets	
	<input type="checkbox"/> 5 mg Tablet	<input type="checkbox"/> Take 1 tablet by mouth 2 times daily	<input type="checkbox"/> 60 Tablets	
		<input type="checkbox"/> Other: _____		
	<input type="checkbox"/> 22 mg XR Tablet	<input type="checkbox"/> Take 1 tablet by mouth once daily	<input type="checkbox"/> 30 Tablets	
<input type="checkbox"/> 11 mg XR Tablet	<input type="checkbox"/> Take 1 tablet by mouth once daily	<input type="checkbox"/> 30 Tablets		
<input type="checkbox"/> Zeposia®	<input type="checkbox"/> Titration Pack (7-day)	<input type="checkbox"/> Titration Dose: 0.23 mg by mouth once daily on day 1-4, 0.46 mg once daily on day 5-7, followed by 0.92 mg once daily thereafter	<input type="checkbox"/> 1 Titration Kit	NA
	<input type="checkbox"/> Titration Pack (28-day)			
	<input type="checkbox"/> 0.92 mg Capsule	<input type="checkbox"/> Take 1 tablet by mouth daily	<input type="checkbox"/> 30 Tablets	

Physician signature required

Product substitution permitted

_____ Date _____

Dispense as written

_____ Date _____

Ancillary supplies and kits will be provided as needed for administration.

The information included in this FAX is intended for the sole use of the individual to whom it is addressed and may contain information that is privileged, confidential and exempt from disclosure under applicable law. If you are not the intended addressee, nor authorized to receive for the intended addressee, you are hereby notified that you may not use, copy, disclose or distribute this information. If you have received this FAX in error, please contact the sender and destroy the entire document.